

Better-behaved GKZ systems and toric mirror symmetry


Zengrui Han
(Joint work with Lev Borisov)

Rutgers, the State University of New Jersey

October 28, 2023

Combinatorial set up

- $N = \mathbb{Z}^d$ is a lattice, C is a polyhedral cone whose ray generators are all of height 1.



- $\text{Spec}(\mathbb{C}[C^\vee \cap N^\vee])$ is a (singular) affine toric Gorenstein variety.
- A regular triangulation Σ of C gives a crepant resolution $\mathbb{P}_\Sigma \rightarrow \text{Spec}(\mathbb{C}[C^\vee \cap N^\vee])$, where \mathbb{P}_Σ is a (non-compact) Calabi-Yau toric orbifold.

Definition of better-behaved GKZ systems

For any lattice point $c \in C \cap N$ we attach a holomorphic function $\Phi_c(x_1, \dots, x_n)$, each variable corresponds to a lattice point (of height 1).

Definition (Borisov-Horja, 2010)

The **better-behaved GKZ system** $bbGKZ(C, 0)$ is given by

$$\frac{\partial}{\partial x_i} \Phi_c = \Phi_{c+v_i}, \text{ for any } i, c \in C$$

and

$$\sum_{i=1}^n \mu(v_i) x_i \frac{\partial}{\partial x_i} \Phi_c + \mu(c) \Phi_c = 0, \text{ for any } \mu \in N^\vee, c \in C$$

Similarly we can define $bbGKZ(C^\circ, 0)$ by considering $c \in C^\circ$ only.

Gamma series solutions

- We can construct series solutions over certain region of convergence from the K-theory of \mathbb{P}_Σ . (Some kind of "central charge")

$$K_0(\mathbb{P}_\Sigma) \otimes \mathbb{C} \xrightarrow{\sim} \{\text{Solutions to } bbGKZ(C^\circ)\}$$

$$K_0^c(\mathbb{P}_\Sigma) \otimes \mathbb{C} \xrightarrow{\sim} \{\text{Solutions to } bbGKZ(C)\}$$

Gamma series solutions

- We can construct series solutions over certain region of convergence from the K-theory of \mathbb{P}_Σ . (Some kind of "central charge")

$$K_0(\mathbb{P}_\Sigma) \otimes \mathbb{C} \xrightarrow{\sim} \{\text{Solutions to } bbGKZ(C^\circ)\}$$

$$K_0^c(\mathbb{P}_\Sigma) \otimes \mathbb{C} \xrightarrow{\sim} \{\text{Solutions to } bbGKZ(C)\}$$

- "A-model pairing":

$$\chi : K_0(\mathbb{P}_\Sigma) \times K_0^c(\mathbb{P}_\Sigma) \rightarrow \mathbb{Z}, (\mathcal{F}, \mathcal{G}) \mapsto \sum_{i=0}^{\infty} (-1)^i \dim \text{Ext}^i(\mathcal{F}, \mathcal{G})$$

Gamma series solutions

- We can construct series solutions over certain region of convergence from the K-theory of \mathbb{P}_Σ . (Some kind of "central charge")

$$K_0(\mathbb{P}_\Sigma) \otimes \mathbb{C} \xrightarrow{\sim} \{\text{Solutions to } bbGKZ(C^\circ)\}$$

$$K_0^c(\mathbb{P}_\Sigma) \otimes \mathbb{C} \xrightarrow{\sim} \{\text{Solutions to } bbGKZ(C)\}$$

- "A-model pairing":

$$\chi : K_0(\mathbb{P}_\Sigma) \times K_0^c(\mathbb{P}_\Sigma) \rightarrow \mathbb{Z}, (\mathcal{F}, \mathcal{G}) \mapsto \sum_{i=0}^{\infty} (-1)^i \dim \text{Ext}^i(\mathcal{F}, \mathcal{G})$$

- Question: What is the pairing on the side of solutions ("B-model pairing")?

B-model pairing

- We find an explicit combinatorial formula for the pairing on the B-side.

B-model pairing

- We find an explicit combinatorial formula for the pairing on the B-side.
- Fix a generic $v \in C^\circ$, define the pairing between solution spaces of $bbGKZ(C, 0)$ and $bbGKZ(C^\circ, 0)$ by the following formula:

$$\langle \Phi, \Psi \rangle_{GKZ} := \sum_{\substack{c \in C, d \in C^\circ \\ I \subseteq \{1, \dots, n\}, |I| = \text{rk } N}} \xi_{c,d,I} \text{Vol}_I \left(\prod_{i \in I} x_i \right) \Phi_c \Psi_d$$

where

$$\xi_{c,d,I} = (-1)^{\deg c}, \text{ if } \dim \sigma(I) = \text{rk } N, c + \epsilon v, d - \epsilon v \in \sigma(I)^\circ \\ \text{otherwise } 0.$$


Theorem (Borisov-H., 2023)

- The B-model pairing $\langle \Phi, \Psi \rangle_{GKZ}$ provides a non-degenerate pairing

$$bbGKZ(C, 0) \times bbGKZ(C^\circ, 0) \rightarrow \mathbb{C}.$$

- For any regular triangulation Σ , the B-model pairing $\langle \Phi, \Psi \rangle_{GKZ}$ agrees with the A-model pairing $\chi : K_0(\mathbb{P}_\Sigma) \times K_0^c(\mathbb{P}_\Sigma) \rightarrow \mathbb{C}$.

Example: $[\mathbb{C}^2/\mathbb{Z}_3]$








- In this case the formula is

$$\begin{aligned} \langle \Phi, \Psi \rangle_{GKZ} &= \Phi_{00}(x_0 x_1 \Psi_{12} + 2x_0 x_2 \Psi_{22} + 3x_0 x_3 \Psi_{32}) \\ &\quad - \Phi_{11}(2x_0 x_2 \Psi_{11} + 3x_0 x_3 \Psi_{21}) - \Phi_{21}(3x_0 x_3 \Psi_{11}) \end{aligned}$$

- Consider e.g. $\mathcal{O}(D_3) \in K_0(\mathbb{P}_\Sigma)$ and $\mathcal{O}_{D_1}(D_3) \in K_0^c(\mathbb{P}_\Sigma)$, they give solutions $\Psi = (\Psi_d)_{d \in C^\circ}$ and $\Phi = (\Phi_c)_{c \in C}$ (rather transcendental).
- Plugging them into the formula above yields

$$\langle \Phi, \Psi \rangle_{GKZ} = \chi(\mathcal{O}(D_3), \mathcal{O}_{D_1}(D_3)) = 1$$

References

-  L. Borisov, R.P. Horja, *On the better behaved version of the GKZ hypergeometric system*. *Mathematische Annalen* 357.2 (2013): 585-603.
-  L. Borisov, R.P. Horja, *Applications of homological mirror symmetry to hypergeometric systems: duality conjectures*. *Advances in Mathematics* 271 (2015): 153–187.
-  L. Borisov, Z. Han, *On hypergeometric duality conjecture*. arXiv:2301.01374.
-  Z. Han, *Analytic continuation of better-behaved GKZ systems and Fourier-Mukai transforms* arXiv: 2305.12241.
-  L. Borisov, Z. Han, C. Wang, *On duality of certain GKZ hypergeometric systems*. *Asian Journal of Mathematics* 25.1 (2021): 65–88.

Thank you!